

DOI: 10.1590/2317-4889202020190120

Mineralogical evolution of the northern Bossoroca ophiolite, São Gabriel terrane
Amanda Juliano Massuda, Léo Afraneo Hartmann¹, Gláucia Nascimento Queiroga,
Marco Paulo de Castro, Carolina Gonçalves Leandro, Jairo Francisco Savian

Supplementary Table 10. Electron microprobe analyses of talc.

Chromite-talc-magnesite granofels, sample C3P10

	29	30	33	34
SiO ₂	61.82	61.14	61.64	62.09
TiO ₂	0.02	0.03	0.04	0.01
Al ₂ O ₃	0.05	0.44	0.15	0.17
Cr ₂ O ₃	0.00	0.03	0.00	0.02
Fe ₂ O ₃	0.00	0.00	0.00	0.00
FeO	1.58	1.63	1.59	1.66
MnO	0.01	0.00	0.00	0.00
MgO	30.47	30.00	30.11	30.38
CaO	0.02	0.03	0.03	0.00
Na ₂ O	0.04	0.14	0.08	0.07
K ₂ O	0.03	0.17	0.04	0.07
Total	94.04	93.61	93.68	94.47
Structural formulae based on 7 oxygen				
Si	2.537	2.525	2.539	2.537
Ti	0.001	0.001	0.001	0.000
Al	0.002	0.021	0.007	0.008
Cr	0.000	0.001	0.000	0.001
Fe (iii)	0.000	0.000	0.000	0.000
Fe (ii)	0.054	0.056	0.055	0.057
Mn	0.000	0.000	0.000	0.000
Mg	1.863	1.847	1.848	1.850
Ca	0.001	0.001	0.001	0.000
Na	0.003	0.011	0.006	0.006
K	0.002	0.009	0.002	0.004
Total	4.463	4.472	4.459	4.463