

DOI: 10.1590/2317-4889202020190120

Mineralogical evolution of the northern Bossoroca ophiolite, São Gabriel terrane
Amanda Juliano Massuda, Léo Afraneo Hartmann¹, Gláucia Nascimento Queiroga,
Marco Paulo de Castro, Carolina Gonçalves Leandro, Jairo Francisco Savian

Supplementary Table 13. Electron microprobe analyses of prehnite. BaO, Rb₂O, Cs₂O, ZnO, F, B₂O₃ below detection limit.

Amphibolite, sample C3P2

Analysis	23	24	25
SiO ₂	42.01	42.84	42.50
TiO ₂	0.03	0.06	0.04
Al ₂ O ₃	23.77	23.47	23.33
FeO	3.41	3.54	3.31
MnO	0.03	0.00	0.05
MgO	1.15	1.49	0.49
CaO	23.90	23.21	24.22
Na ₂ O	0.09	0.02	0.04
K ₂ O	0.20	0.16	0.05
Cl	0.074	0.171	0.01
H ₂ O *	4.25	4.25	4.25
O=F,Cl	0.02	0.04	0.00
Total	98.89	99.17	98.28
Structural formulae based on 22 O			
Si	5.905	5.986	5.999
Al total	3.938	3.866	3.881
Ti	0.003	0.006	0.004
Fe	0.401	0.414	0.391
Mn	0.004	0.000	0.006
Mg	0.241	0.310	0.103
Zn	0.000	0.000	0.000
Ca	3.600	3.475	3.663
Na	0.025	0.005	0.011
K	0.036	0.029	0.009
OH *	3.982	3.959	3.998
Cl	0.018	0.041	0.002
Total	18.153	18.092	18.067